

Reg. No.: 1471/2009 M.O.H



AMOXYVETo-15

Injectable Suspension

Composition

Each 1 ml contains:
Amoxicillin Trihydrate 172.2 gm
(equivalent to Amoxicillin base 150 mg)

Indications

Bacterial infections caused by microorganisms susceptible to penicillins, such as infections of the alimentary tract (bacterial gastroenteritis), respiratory tract (tracheobronchitis), urogenital tract (cystitis) & colimastitis.

Cattle, Calves Clostridium, A. pyogenes, Corynebacterium sp., Anthrax, Streptococcal mastitis, listeriosis, infectious keratoconjunctivitis.



Dosage and administration

I/M or S/C injection.

For active ingredient

Cattle, calves: 10 mg amoxicillin/kg B.W.

For whole product

Cattle, calves: 1 ml/15 kg B.W.

Frequency of injection:

one injection to be repeated after 48 hours.

Withdrawal time

Meat: 28 days. Milk: 2 days.

Storage

Store between 2°C and 25 °C. use within 1 months from first use.

Warning and precaution

Avoid the use of quick working bacteriostatic medicines, like tetracyclines and chloramphenicol when using this medicine. Shake well before use to obtain homogenate suspension.

Package

100 ml glass injection bottles.

FOR VETERINARY USE ONLY

Manufactured by Laboratoires Biové France for V.M.D, n.v. Belgium

Exclusive agent: INTERMEDICAVET, Animal Health products.

Distributed by: WAYZON Pharma.