

Reg. No.: 3896/2016 M.O.H



CYDECTIN TRICLAMOX

Oral Solution for sheep

Composition

Each 1 ml contains:

Moxidectin	1 mg.
Triclabendazole	50 mg.

Properties:

Moxidectin is an endectocide active against a wide range of internal and external parasites and is a second-generation macrocyclic lactone of the milbemycin family. Moxidectin interacts with GABA receptors and chloride channels. The net effect is to open the chloride channels on the postsynaptic junction to allow the inflow of chloride ions and induce an irreversible resting state. This results in flaccid paralysis and the eventual death of parasites exposed to the drug.

Triclabendazole is a flukicide belonging to the benzimidazole group of anthelmintics. It is well established that benzimidazole anthelmintics selectively bind to β -tubulin, thus causing the depolymerization of microtubules and the subsequent disruption of microtubule-based processes in helminths.

Indications

For the treatment of mixed nematode and fluke infections, caused by moxidectin and triclabendazole-sensitive strains.

Gastro-intestinal nematodes:

Haemonchus contortus, Teladorsagia circumcincta, Ostertagia trifurcata, Trichostrongylus axei, Trichostrongylus colubriformis, Trichostrongylus vitrinus, Nematodirus battus, Nematodirus spathiger, Nematodirus filicollis, Strongyloides papillosus, Cooperia curticei, Cooperia oncophora, Oesophagostomum columbianum, Oesophagostomum venulosum, Chabertia ovina, Trichuris ovis.

Respiratory tract nematode:

Dictyocaulus filaria.

TREMATODES

Fasciola hepatica: Adult, Late immature, and early immature.



Reg. No.: 3896/2016 M.O.H

Dosage and route of administration

Should be given as a single oral drench of 1 ml/5 kg bodyweight, equivalent to 0.2 mg moxidectin/kg bodyweight and 10 mg triclabendazole/kg body weight, using any standard drenching equipment.

Withdrawal time:

Meat and offal: 31 days.

Milk: not authorized for use in ewes producing milk intended for human consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for human consumption.

Shelf life

Do not store above 25°C. Protect from light, and don't freeze. Shelf-life of the veterinary medicinal product as packaged for sale: 2 years. Shelf-life after first opening the immediate packaging: 6 months

Package

1 liter polyethylene containers with polypropylene screw-cap.

FOR ANIMAL TREATMENT ONLY

zoetis



Licenser Holder: Zoetis Spain.

Manufactured by: Zoetis Spain.

Imported by: Zoetis Egypt.

Exclusive Distributor: WAYZON Pharma.