



Reg. No.: 3168/2008 M.O.H

# MELOXIDEL

Injectable Solution

## Composition

Each 1 ml contains:

**Meloxicam** 5 mg

## Properties

Meloxicam is a Cox2-receptor preferential NSAIDS. Meloxicam has anti-inflammatory, analgesic and antipyretic activity similar to other NSAIDS.

## Mode of action

Meloxidel 0.5% acts by inhibition of cyclooxygenase, phospholipase A2 and inhibition of prostaglandin synthesis.

## Indications

### In Cattle

1. Acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle.
2. Use in diarrhea in combination with oral rehydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle.
3. For the relief of post-operative pain following dehorning in calves.
4. Supportive therapy in the treatment of acute mastitis.

### In Dogs and Cats

1. The alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders.
2. Control of perioperative pain and inflammation following orthopedic and soft tissue surgery.



## Dosage and route of administration

By S/C or I/V injection

For Active Ingredient:

**Cattle** 0.5 mg meloxicam/Kg b.wt. as a single dose.

**Dogs** 0.2 mg meloxicam/Kg b.wt. as a single dose.

**Cats** 0.3 mg meloxicam/Kg b. wt. as a single dose.

For Whole Product:

**Cattle** 10 ml/100 kg b. wt. as a single dose by S/C or I/V injection.

**Dogs** 0.4 ml/10 kg b. wt. as a single dose by S/C injection.

**Cats** 0.3 ml / 5 kg b. wt. as a single dose by S/C injection.

## Use during pregnancy

**Cattle** can be used during pregnancy.

## Warning and precautions

1. Not to be used in animals less than 6 weeks of age or aged animals.
2. Not to be used in dehydrated and hypovolemic or hypotensive animals.
3. Not to be used for breeding, pregnant or lactating dogs.
4. Concurrent antibacterial should be given if bacterial infection is present.

## Contra-indications

1. Should not to be used for animals with cardiac, hepatic or renal disease where the possibility of gastro-intestinal ulceration or bleeding.
2. Should not to be used for animals with hypersensitivity to meloxicam.
3. Not to be used for animals during treatment with other NSAIDS, Corticosteroids or anticoagulants concurrently or within 24 hours.
4. Not to be used for cats less than 2 kg body weight.
5. Do not use in animals of less than one week of age.

## Drug interactions

1. Avoid concurrent administration of potentially nephrotoxic drugs, anticoagulants and aminoglycosides.
2. Not be used for plasma-protein bound drugs as meloxicam is highly bound to plasma proteins it can displace or be displaced by other highly protein bound drugs including warfarin and phenylbutazone.

Reg. No.: 3168/2008 M.O.H

**MELOXIDEL**  
Injectable solution

## Side effects

Prolonged use may cause gastro-intestinal lesions, Inappetance, vomiting, diarrhea, transient swelling at injection site.

## Withdrawal time

**Cattle** Meat and offal: 15 days.  
Milk: 5 days.

## Package

30, 50 and 100 ml in a glass vial with outer label in carton box and insert leaflet.

**FOR VETERINARY USE ONLY**



PRODUCED BY DELTA PHARMACEUTICAL INDUSTRIES

EXCLUSIVE DISTRIBUTOR: WAYZON PHARMA