

Reg. No.: 3280/2010 M.O.H



# TYLOVETo-20

Injectable Solution

## Composition

Each 1 ml contains:  
Tylosine base (as tartrate) 200 mg.

## Indications

Infection caused by micro organisms susceptible to tylosin, such as respiratory infections (pneumonia,...). Dysentery and arthritis caused by mycoplasmas, mastitis and endometritis.



## Dosage and administration

Administer TYLOVETo-20 by deep intramuscular injection.

### For active ingredient

Cattle: 5-10 mg tylosin/kg B.W.

### For whole product

Cattle: 0.5-1 ml/20 kg B.W.

Inject every 12 hours.

Do not administer for more than 5 consecutive days.

## Withdrawal time

Meat: 21 days.      Milk: 4 days.

## Storage

Store at temperature not exceeding 25°C and protected from light. Shelf life of the product as packaged for sale 2 years. Shelf life after first opening 14 days.

## Adverse effects

Due to the irritant nature of macrolides, phlebitis may occur after intravenous administration of the product. Local intolerance (swelling and edema) at the injection site was noted in 80-90% of the treated animals, this reaction gradually diminishes after 5 days.

## Contraindications

Do not inject in horses. Do not inject in animals with known hypersensitivity to macrolides (erythromycin, spiramycin,...).

## Package

100 ml glass injection bottles.

## FOR VETERINARY USE ONLY

Manufactured by Laboratoires Biové France for V.M.D, n.v. Belgium

**Exclusive agent:** INTERMEDICAVET, Animal Health products.

**Distributed by:** WAYZON Pharma.